

Tobacco

Any Current Tobacco Use

Any current tobacco use is adults reporting current smoking or smokeless tobacco use. **31.2%** of adults report any current tobacco use.

Cigarette Smoking

Cigarette smoking is an adult reporting having smoked at least 100 cigarettes in their lifetime and are currently smoking everyday or some days. **25.8%** of adults report cigarette smoking.


Daily Cigarette Smoking

Daily cigarette smoking is an adult reporting currently smoking cigarettes every day. **19.5%** of adults report daily cigarette smoking.

Smokeless tobacco Use for Males


Smokeless tobacco use for males is men reporting currently using smokeless tobacco products such as chewing tobacco or snuff. **14.7%** of adult men report smokeless tobacco use.

Tobacco Use in Natrona County, 2011-2015



Source: Wyoming BRFSS


Chronic Lower Respiratory Disease Mortality Rate



- Natrona has a higher mortality rate for Chronic Lower Respiratory Disease than Wyoming and The United States
- From 2011 to 2015, Natrona had a mortality rate of **78.7** per 100,000 for Chronic Lower Respiratory Disease

Source: US Department of Health and Human Services

Adolescent Cigarette Use



Adolescent Tobacco Use

Cigarette Use


- Cigarette use for adolescents is adolescents reporting cigarette use in the last 30 days
- There has been a decline in cigarette use for 12th graders and 8th graders. Cigarette use for these grades are at the lowest
- **12%** of 12th graders report cigarette use in 2016
- In 2016, **4%** of 8th graders report cigarette use
- Cigarette use has risen since 2014 for 6th graders and 10th graders
- **2%** of 6th graders report cigarette use in 2016
- In 2016, **11%** of 10th graders reported cigarette use

Tobacco Use and Pregnancy

Source: PNA Survey

- Women who report smoking during pregnancy
- Women who quit smoking during pregnancy
- Women who quit smoking before pregnancy

Pregnant Women Reporting Smoking



Source: Wyoming Department of Health Chronic Disease and Maternal Child Health

- Natrona has a higher rate of women who report smoking during pregnancy than Wyoming.
- **30%** of women surveyed reported quitting smoking before pregnancy
- In 2016, **27.1%** of women surveyed reported quitting smoking during pregnancy
- The amount of women who reported smoking during pregnancy decreased from 2014 at **20.5%** to **15.2%** in 2016